


Significance of Isolated Non-Prompt Photons on Photon-Triggered Jet Observables

Chathuranga Sirimanna
Duke University

Workshop on Heavy Ion Physics in the EIC Era

Outline

- Introduction: JETSCAPE framework and multistage evolution
- Photon triggered jets
- Simulating jet evolution with JETSCAPE framework
- Simultaneous description of leading hadron and jet spectrum
- This study: 5.02 TeV
 - ❖ γ -jet: asymmetry, correlation, and R_{AA}
 - ❖ $\gamma + 2$ jets
 - ❖ Groomed jets substructure using photon triggered jets
- Summary



<https://github.com/JETSCAPE/JETSCAPE>




Manual ([arXiv:1903.07706](https://arxiv.org/abs/1903.07706)), JETSCAPE PP19 tune ([arXiv:1910.05481](https://arxiv.org/abs/1910.05481)), JETSCAPE AA23 tune ([arXiv:2204.01163](https://arxiv.org/abs/2204.01163))

Duke

Introduction

- **JETSCAPE**: General, modular and extensive framework
- No single model can describe all stages of jet evolution
- Multi-stage jet evolution
 - ❖ Different stages depending on the virtuality, Q and Energy, E of the partons
- One can customize the framework by using their own modules
- ASCII, Gzip, and HepMC output formats



Pre-generated Trento Initial

Pre generated 2+1D VISHNU
Hydro profiles

Pythia Hard Scattering


MATTER – Medium modified DGLAP evolution in High
Virtuality

LBT – Linearized Boltzmann Transport model with HT
gluon emission

Duke

JETSCAPE framework: Multistage Evolution

- Large Q , Large E : Dominated by radiation with few scatterings (DGLAP, HT)
 - ❖ **MATTER** (*Majumder(13), Kordell, Majumder(17), Cao, Majumder(17)*)
- Small Q , Large E : Scattering driven emission, mostly by medium effects (Transport, AMY, HT)
 - ❖ **LBT** (*Wang, Zhu(13), Luo, et al.(15,18), Cao, et al.(16,17), He, et al.(18)*)
 - ❖ **MARTINI** (*Schenke, Gale, Jeon(09), Park, Jeon, Gale(17, 18)*)
- Small Q & E : Nearly thermal, strongly coupled (AdS/CFT)
 - ❖ **AdS/CFT** (*Chesler, Rajagopal(14, 15), Pablos, et al.(15, 16, 17), and others*)



Virtuality Separation Scale: Q_0

Switching between modules parton by parton depending on the virtuality and energy



Large Q : $Q > Q_0$

Small Q : $Q < Q_0$

Duke

Prompt photons as Probes of QGP

- Prompt photons - produced directly in the hard sub-processes
- Can be used to estimate the energy and the direction of jet initiating parton (before the energy loss) – Calibrated probe of the QGP
 - ❖ Limited Statistics: Challenging to measure experimentally
- Important probe to study jet energy loss (Wang, Huang, and Sarcevic, PRL 77 (1996) 231-234)
- Isolation criteria is necessary to identify the prompt photons
 - ❖ Same isolation criteria used in experimental analysis (CMS-HIN-13-006, CMS-HIN-16-002, PLB 789 (2019) 167)
- Isolated photons mainly consist of prompt photons
 - ❖ Isolated Non-prompt photons make considerable contribution



Simulating Jet Evolution



➤ **p-p baseline:** Pythia for hard scattering and MATTER for shower

➤ **PbPb:**

- ❖ Virtuality separation scale, $Q_0 = 2 \text{ GeV}$
- ❖ MATTER+LBT with recoil
- ❖ Colorless hadronization (color is randomly assigned) with Pythia Lund String model
- ❖ Event by event hydro


➤ **p-p and PbPb:**

- ❖ Prompt photons, photons from intermediate shower and fragmentation photons
- ❖ Same isolation criteria used in experimental analysis



$$\sum_{\Delta R} E_T < E_{T, Iso}$$

$$\Delta\phi_{\gamma, jet} < \frac{7\pi}{8}$$



Simultaneous Description of Leading Hadrons and Jets

➤ Parameters tuned for simultaneously describe leading hadron and jet spectra

❖ Blue line of each plot: $Q_{sw} = Q_0 = 2 \text{ GeV}$, $\alpha_s^{fix} = 0.3$, $\tau_0 = 0.6 \text{ fm}/c$, and $T_c = 160 \text{ MeV}$



❖ AA23 tune

➤ Not tuned using Bayesian calibration

➤ Same tune can be used to accurately describe number of different observables (Different E_{CM} , centrality, etc.)

➤ JETSCAPE AA paper: Phys.Rev.C 107 (2023) 3, 034911, arXiv: [2204.01163](https://arxiv.org/abs/2204.01163)


❖ Further information on parameter tuning



γ -jet Asymmetry – p-p

- γ -jet Asymmetry: $X_{J\gamma} = \frac{p_T^{jet}}{p_T^\gamma}$
- 5.02 TeV p-p: Full Events and Prompt Photon Events
 - ❖ $p_T^{jet} > 31.6 \text{ GeV}$; $|\eta_\gamma| < 2.37$ (excluding the region $1.37 < |\eta_\gamma| < 1.52$); $R = 0.4, |\eta_{jet}| < 2.8, |\Delta\phi| > \frac{7\pi}{8}$
 - ❖ Isolation cut ($E < 3 \text{ GeV}$) $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.3$
- p-p simulation using JETSCAPE PP19 tune
- Full events: Have better description with relatively large error bars

ATLAS [PLB 789, 167-190 (2019)]



γ -jet Asymmetry – Central PbPb


➤ 5.02 TeV PbPb: Full Events and Prompt Photon Events

- ❖ $p_T^{jet} > 31.6 \text{ GeV}$; $|\eta_\gamma| < 2.37$ (excluding the region $1.37 < |\eta_\gamma| < 1.52$); $R = 0.4, |\eta_{Jet}| < 2.8, |\Delta\phi| > \frac{7\pi}{8}$
- ❖ Isolation cut ($E < 8 \text{ GeV}$) $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.3$

➤ Unfolded experimental results


➤ Full events: Have better description with relatively large error bars

ATLAS [PLB 789, 167-190 (2019)]






γ -jet Asymmetry – p-p and Central PbPb

- 5.02 TeV: Full Events and Prompt Photon Events
 - ❖ $p_T^{jet} > 30 \text{ GeV}$, $|\eta_\gamma| < 1.44$, $R = 0.3$, $|\eta_{jet}| < 1.6$, $|\Delta\phi| > \frac{7\pi}{8}$
 - ❖ Isolation cut ($E < 5 \text{ GeV}$) $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.4$
- Full events: Have better description with large error bars
- Same JETSCAPE Events
- Smeared jet p_T (p-p and PbPb)



CMS [PLB 785, 14-39 (2018)]

Smearing
Function



γ -jet Asymmetry – Smearred p-p

➤ 5.02 TeV p-p: Full Events and Prompt Photon Events


- ❖ $p_T^{jet} > 30 \text{ GeV}$, $|\eta_\gamma| < 1.44$, $R = 0.3$, $|\eta_{jet}| < 1.6$, $|\Delta\phi| > \frac{7\pi}{8}$
- ❖ Isolation cut ($E < 5 \text{ GeV}$) $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.4$

➤ Ratio plots: Shows large deviation at large $X_{J\gamma}$

- ❖ Wide angle photon radiation after initial hard scattering

➤ Full events: Have better description with large error bars

CMS [PLB 785, 14-39 (2018)]



γ -jet Asymmetry – Peripheral PbPb

- 5.02 TeV PbPb: Full Events and Prompt Photon Events
- Deviated from experimental results
 - ❖ Similar behavior can be seen in the central events
- Full Events: Significantly better agreement
 - ❖ Larger Statistical Errors
 - ❖ Need larger number of events: More computer power
- Non-prompt photons: Important for direct photon observables



ATLAS [PLB 789, 167-190 (2019)]

γ -jet Correlation - pp


➤ 5.02 TeV pp: Full Events and Prompt Photon Events

➤ Prompt Photon Events: Deviated from experimental results

➤ Full Events: Significantly better agreement

- ❖ Larger Statistical Errors
- ❖ Overestimate the smallest bin

➤ $\Delta\phi > \frac{7\pi}{8}$: Similar in large p_T^γ



CMS [PLB 785, 14-39 (2018)]

γ -jet Correlation - PbPb

➤ 5.02 TeV PbPb: Full Events and Prompt Photon Events

➤ Same Behavior as pp


➤ Prompt Photon Events: Deviated from experimental results

➤ Full Events: Significantly better agreement

❖ Larger Statistical Errors

❖ Overestimate the smallest bin


➤ $\Delta\phi > \frac{7\pi}{8}$: Similar in large p_T^γ



CMS [PLB 785, 14-39 (2018)]

γ -jet- R_{AA}

- 5.02 TeV: Full Events and Prompt Photon Events
- Two centralities
- Full events captures most of the features
- Statistical Errors are significant
- More statistics or Different approach?
- Full Events: Significantly better agreement
 - ❖ Larger Statistical Errors
 - ❖ Need larger number of events: More computer power
- Non-prompt photons: Important for direct photon observables







ATLAS [PLB 846, 138154 (2023)]

$\gamma+2$ jets


- Same Isolation criteria as γ -triggered single jet analysis
- Prompt Photon Events only
- $PbPb/pp$ ratio for all three observables have good agreement
- Multiplicative factor of 1.4 separately improves Pb-Pb and p-p
- Tension with models (without multiplicative factor)
- Possible role of NLO effects

ATLAS Preliminary [ATLAS-CONF-2023-008]



Groomed Jet Substructure with γ -triggered jets



- Prompt Photon Events Only
- Soft Drop grooming ($z_{cut} = 0.2, \beta = 0$)
- Prominent modification of quark jets
- Manifestation of quark jet characteristics in γ -tagged jets



$$m_g = \sqrt{(p_1 + p_2)_\mu (p_1 + p_2)^\mu}$$

r_g -distribution



m_g -distribution




Groomed Jet Substructure with γ -triggered jets

- z_g : energy imbalance of its hardest splitting
- r_g : angular separation of its hardest splitting
- Doesn't show significant dependence on $X_{J\gamma}$
- Only a slight signal of narrowing of the splitting can be seen in r_g

$$z_g = \frac{\min(p_{T,i}, p_{T,j})}{p_{T,i} + p_{T,j}}$$



Unlocking the Power of Multistage Evolution



Summary and Future Directions

- Photon observables - an independent, parameter free verification of the multistage evolution
- Using the prompt photon events from hard scattering might not be sufficient
 - ❖ Prompt photon events are rare: Computer intensive simulation
 - ❖ Full events shows a better description of all results with relatively large error bars
 - ❖ Isolated Non-prompt photons make considerable contribution
- Further studies needed to understand $\gamma+2$ jets
- Groomed Jet substructure doesn't show significant dependence to $X_{J\gamma}$
- Full event analysis with reduced statistical uncertainty for both central and peripheral PbPb events
 - ❖ More statistics or Different approach?
- Include more physics in our simulations