

[252/151]

INT WORKSHOP INT-20R-2C

# Accessing and Understanding the QCD Spectra

March 20, 2023 - March 24, 2023

Mar, 2023

# CAN CONSTITUENT GLUONS DESCRIBE GLUEBALLS AND HYBRIDS?

Eric Swanson



# References

Light hybrid decays,  
C. Farina & E.S. Swanson, in preparation.

Light hybrid mixing and phenomenology,  
E.S. Swanson, arXiv:2302.01372.

Heavy hybrid decays in a constituent gluon model,  
C. Farina, H.G. Tecocoatzi, A. Giachino, E. Santopinto, & E.S. Swanson,  
*Phys.Rev.D* 102 (2020) 1, 014023.

The low lying glueball spectrum,  
A.P. Szczepaniak & E.S. Swanson, *Phys.Lett.B* 577 (2003) 61–66.

Coulomb gauge QCD, confinement, and the constituent representation,  
A.P. Szczepaniak & E.S. Swanson, *Phys.Rev.D* 65 (2001) 025012.



Christian Farina

# Flavour *Mixing* in the Isoscalar Sector a la LGT





FIG. 11: Isoscalar (green/black) and isovector (blue) meson spectrum on the  $m_\pi = 391$  MeV,  $24^3 \times 128$  lattice. The vertical height of each box indicates the statistical uncertainty on the mass determination. States outlined in orange are the lowest-lying states having dominant overlap with operators featuring a chromomagnetic construction – their interpretation as the lightest hybrid meson supermultiplet will be discussed later.


# Hybrid Flavour Mixing is Different!

## ISOSCALARS



+  $U(2)_W$   
Anomaly  
(=  $U(1)$ ?)


+ centonum  
mixing!






$\sim \alpha_s^n (40)^2$

or  $U(1)_B \times U(1)_L$

$g^+ / g^b$



## HYBRIDS



We need a specific model for hybrids and glueballs, interactions, etc

The form of the gluonic structure present in the operators having good overlap with these states is chromomagnetic, having  $J_g^{P_g C_g} = 1^{+-}$ . With the  $q\bar{q}$  pair in an internal  $S$ -wave this describes the observed  $J^{PC}$ . Heav-



The lightest hybrid meson supermultiplet in QCD  
J.J. Dudek, Phys.Rev.D 84 (2011) 074023

Agrees with old bag models and other modelling


Heavy hybrids with constituent gluons,  
E.S. Swanson & A.P. Szczepaniak, Phys.Rev.D 59 (1999) 014035

# Glueballs


# Glueballs



$V_{4g}$




$H_{ggg}$




$K^{(1)}$

$c(\Lambda)$



$\Delta E < \Lambda$



$\Delta E > \Lambda$

$$|JM; \lambda, \lambda'\rangle = \frac{1}{\sqrt{2(N_c^2 - 1)}} \sqrt{\frac{2J+1}{4\pi}} \int \frac{d^3k}{(2\pi)^3} \psi(k) D_{M, \lambda-\lambda'}^{J*}(\phi, \theta, -\phi) \Pi a^\dagger(k, \lambda, A) a^\dagger(-k, \lambda, A) |0\rangle$$

$$|JM; \eta\rangle = \frac{1}{\sqrt{2}} (|JM; \lambda, \lambda'\rangle + \eta |JM; -\lambda, -\lambda'\rangle)$$



# Glueballs

$$E \int \frac{k^2 dk}{(2\pi)^3} |\psi_i(k)|^2 = \int \frac{k^2 dk}{(2\pi)^3} 2\omega(k) |\psi_i(k)|^2 + \frac{N_C}{2} \sum_i \int \frac{k^2 dk}{(2\pi)^3} \frac{q^2 dq}{(2\pi)^3} \frac{\omega(k)}{\omega(q)} \left[ \frac{4}{3} V_0 + \frac{2}{3} V_2 \right] |\psi_i(k)|^2$$

$$- \frac{N_C}{4} \int \frac{k^2 dk}{(2\pi)^3} \frac{q^2 dq}{(2\pi)^3} \frac{(\omega(k) + \omega(q))^2}{\omega(k)\omega(q)} \psi_i^*(q) K_{ij}(q, k) \psi_j(k)$$

$J^P = (\text{odd} \geq 3)^+$  (there is no  $1^+$  gg glueball):

$$K = \frac{J+2}{2J+1} V_{J-1} + \frac{J-1}{2J+1} V_{J+1};$$


$J^P = (\text{even} \geq 0)^-$ :

$$K = \frac{J}{2J+1} V_{J-1} + \frac{J+1}{2J+1} V_{J+1};$$

$J^P = 0^+$ :


$$K = \frac{2}{3} \left( V_0 + \frac{V_2}{2} \right).$$

# Glueballs



# Hybrids

# Hybrids



$$\begin{aligned}
 |JM[LS\ell j_g \xi]\rangle &= \frac{1}{2} T_{ij}^A \int \frac{d^3 q}{(2\pi)^3} \frac{d^3 k}{(2\pi)^3} \Psi_{j_g; \ell m_\ell}(\mathbf{k}, \mathbf{q}) \sqrt{\frac{2j_g + 1}{4\pi}} D_{m_g \mu}^{j_g*}(\hat{k}) \chi_{\mu, \lambda}^{(\xi)} \\
 &\times \langle \frac{1}{2} m \frac{1}{2} \bar{m} | SM_S \rangle \langle \ell m_\ell, j_g m_g | LM_L \rangle \langle SM_S, LM_L | JM \rangle b_{\mathbf{q} - \frac{\mathbf{k}}{2}, i, m}^\dagger d_{-\mathbf{q} - \frac{\mathbf{k}}{2}, j, \bar{m}}^\dagger a_{\mathbf{k}, A, \lambda}^\dagger | 0 \rangle.
 \end{aligned}$$

# Hybrids

TABLE II:  $J^{PC}$  Hybrid Multiplets.

multiplet	operator	$\xi$	$j_g$	$\ell$	$L$	$J^{PC}$ $S = 0$ ( $S = 1$ )
$H_1$	$\psi^\dagger \mathbf{B} \chi$	-1	1	0	1	$1^{--}, (0, 1, 2)^{-+}$
$H_2$	$\psi^\dagger \nabla \times \mathbf{B} \chi$	-1	1	1	1	$1^{++}, (0, 1, 2)^{+-}$
$H_3$	$\psi^\dagger \nabla \cdot \mathbf{B} \chi$	-1	1	1	0	$0^{++}, (1^{+-})$
$H_4$	$\psi^\dagger [\nabla \mathbf{B}]_2 \chi$	-1	1	1	2	$2^{++}, (1, 2, 3)^{+-}$

# Hybrids

Hartree-Fock-type method


$$\Psi_{j_g; \ell m_\ell}(\mathbf{k}, \mathbf{q}) = \chi_{j_g}(k) \varphi_\ell(q) Y_{\ell, m_\ell}(\hat{q}).$$

$$K_q \varphi + \int \chi^* K_g \chi \cdot \varphi + \int \chi^* V \chi \cdot \varphi = E \varphi$$


$$K_g \chi + \int \varphi^* K_q \varphi \cdot \chi + \int \varphi^* V \varphi \cdot \chi = E \chi$$

# Hybrids

Hartree-Fock-type method




# Hybrids





# Hybrids



# Hybrid Flavour Mixing


# Mixing

$$\begin{array}{cccc}
 & |u\bar{u}\rangle & |d\bar{d}\rangle & |s\bar{s}\rangle & |gg\rangle \\
 H_{uds} = & \begin{pmatrix}
 m + A_{nn} & A_{nn} & A_{ns} & \mathcal{A}_n^{(0)} \\
 A_{nn} & m + A_{nn} & A_{ns} & \mathcal{A}_n^{(0)} \\
 A_{ns} & A_{ss} & m + \Delta m + A_{ss} & \mathcal{A}_s^{(0)} \\
 \mathcal{A}_n^{(0)} & \mathcal{A}_n^{(0)} & \mathcal{A}_s^{(0)} & M_{gb}
 \end{pmatrix}
 \end{array}
 \cdot
 \begin{array}{c}
 \\
 \\
 \\
 \end{array}
 \begin{array}{c}
 \\
 \\
 \text{☞} \\
 \end{array}$$


$$H_{iso} = \begin{pmatrix}
 m & 0 & 0 & 0 \\
 0 & m + 2A_{nn} & \sqrt{2}A_{ns} & \sqrt{2}\mathcal{A}_n^{(0)} \\
 0 & \sqrt{2}A_{ns} & m + \Delta m + A_{ss} & \mathcal{A}_s^{(0)} \\
 0 & \sqrt{2}\mathcal{A}_n^{(0)} & \mathcal{A}_s^{(0)} & M_{gb}
 \end{pmatrix}
 \cdot$$

truncate sums over glueballs

# Mixing



$\ell = 0; S = 1; H_1$  only



$$A_{ff'} = \frac{1}{m_f m_{f'}} \int \frac{k^2 dk}{(2\pi)^3} \frac{d^3 q}{(2\pi)^3} \frac{d^3 q'}{(2\pi)^3} \Psi_f(k, \mathbf{q}) \Psi_{f'}^*(k, \mathbf{q}') k^2 V(k) B_J$$


$$= \frac{F_f F_{f'}}{8 m_f m_{f'}} \int \frac{k^2 dk}{(2\pi)^3} |\chi_1(k)|^2 k^2 V(k) B_J.$$

$$\mathcal{A}_f^{(n)} = -\frac{i[gF_f]}{4} \int \frac{k^2 dk}{(2\pi)^3} \frac{\psi_n^*(k) \chi(k)}{\sqrt{\omega(k)}} C_J,$$


$$[gF_f] = \int \frac{d^3 q}{(2\pi)^3} \sqrt{4\pi\alpha_V(q)} \phi_{\ell=0}(q)$$

"octet decay constant, F"

# Mixing



# Mixing



# Mixing


$J^{PC}$	nominal state	$u\bar{u}g$	$s\bar{s}g$	$gg$
$1^{--}$	light	$\approx 100$	$\approx 0$	$\approx 0$
	heavy	$\approx 0$	$\approx 100$	$\approx 0$
	glueball	$\approx 0$	$\approx 0$	$\approx 100$
$0^{-+}$	light	62 [87]	17 [6]	21 [7]
	heavy	24 [8]	75 [91]	0.5 [1]
	glueball	13 [5]	8 [3]	78 [92]
$1^{-+}$	light	37	63	0
	heavy	63	37	0
	glueball	0	0	100
$2^{-+}$	light	54 [59]	46 [41]	0.1 [0]
	heavy	32 [38]	67 [61]	1 [1]
	glueball	2 [0.4]	1 [0.6]	97 [99]

LGT assumptions are validated

# Vector-Hybrid Vector Mixing



# Vector-Hybrid Vector Mixing



$$\mathcal{H} \equiv \langle q\bar{q}g | ig \int \psi^\dagger \alpha \cdot A \psi | q\bar{q} \rangle$$

$$\mathcal{H}_n = -i \frac{g}{m} \frac{2\sqrt{4\pi}}{3} \int \frac{d^3q}{(2\pi)^3} \frac{k^2 dk}{(2\pi)^3} \frac{k}{\sqrt{\omega(k)}} \Psi^*(k, q) \psi_n(q + k/2).$$

$$\mathcal{H}_{1S} = -ig \begin{cases} 84 \text{ MeV}^2/m_q, & \rho \\ 190 \text{ MeV}^2/m_c, & J/\psi \\ 225 \text{ MeV}^2/m_b, & \Upsilon \end{cases} \approx -i \begin{cases} 210 \text{ MeV}, & \rho \\ 60 \text{ MeV}, & J/\psi \\ 20 \text{ MeV}, & \Upsilon \end{cases}$$


Hybrid configuration content of heavy S-wave mesons,  
 [MILC] T. Burch & D. Toussaint, Phys. Rev. 68, 094504 (2003).

$$\mathcal{H}_{NRQCD} \approx 170 \text{ MeV} (J/\psi)$$

$$\mathcal{H}_{NRQCD} \approx 70 \text{ MeV} (\Upsilon).$$


# Vector Hybrid Production/Decay Constant

# Vector Hybrid Production/Decay Constant



$$\delta H = \begin{pmatrix} m_1 & & & & & & \mathcal{H}_{1S} \\ & m_2 & & & & & \mathcal{H}_{2S} \\ & & m_3 & & & & \mathcal{H}_{3S} \\ & & & m_4 & & & \mathcal{H}_{4S} \\ & & & & m_5 & & \mathcal{H}_{5S} \\ & & & & & m_6 & \mathcal{H}_{6S} \\ \mathcal{H}_{1S} & \mathcal{H}_{2S} & \mathcal{H}_{3S} & \mathcal{H}_{4S} & \mathcal{H}_{5S} & \mathcal{H}_{6S} & M_H \end{pmatrix} \cdot$$

# Vector Hybrid Production/Decay Constant



## Vector Hybrid Production/Decay Constant

$$f_H = \frac{1}{\sqrt{M_H}} \sum_{n \neq H} \sqrt{M_n} f_V^{(n)} C_n \quad C_n = \langle nS | H_1(1^{--}) \rangle$$


$$f_V^{(n)} = \sqrt{\frac{3}{M_n}} \int \frac{d^3k}{(2\pi)^3} \psi^{(n)}(\vec{k}) \sqrt{1 + \frac{m_q}{E_k}} \sqrt{1 + \frac{m_{\bar{q}}}{E_{\bar{k}}}} \left( 1 + \frac{k^2}{3(E_k + m_q)(E_{\bar{k}} + m_{\bar{q}})} \right).$$

$$f_{H_1(1^{--})} \approx 20 \text{ MeV}.$$

# Phenomenology


# Phenomenology

lattice hybrid masses vs quark mass



# Phenomenology


lattice hybrid masses vs quark mass






# Phenomenology

lattice hybrid masses vs quark mass




# Phenomenology

lattice hybrid masses vs quark mass




# Phenomenology

lattice hybrid masses vs quark mass



# Phenomenology


lattice hybrid masses vs quark mass



# Phenomenology

TABLE IV: Model Assignments and Experimental Isovector Vector States (MeV).

state	Ref.	mass	width	model (iv)	mass	model (v)	mass
$\rho(770)$	PDG	$775.2 \pm 0.2$	$147.4 \pm 0.8$	$1^3S_1$	720	$1^3S_1$	810
$\rho(1450)$	PDG	$1465 \pm 25$	$400 \pm 60$	$2^3S_1$	1440	$2^3S_1$	1405
$\rho(1570)$	PDG	$1570 \pm 70$	$144 \pm 90$	$1^3D_1$	1510	$1^3D_1$	1497
$\rho(1700)$	PDG	$1720 \pm 20$	$250 \pm 100$	$H_1(1^{--})$	1760	$3^3S_1$	1770
						$2^3D_1$	1835
				$3^3S_1$	1850		
$\rho(1900)$	[32]	$1900 \pm 30$	$50 \pm 30$	$2^3D_1$	1910	$4^3S_1$	2080
$\rho(2150)$	[33]	$2034 \pm 16$	$234 \pm 39$	$4^3S_1$	2170	$H_1(1^{--})$	2100
				$3^3D_1$	2220	$3^3D_1$	2130



# Conclusions

# Conclusions

- a reasonable approximation to lattice glueball and hybrid spectra is obtained (clearly room for improvement)
- hybrid flavour mixing is roughly reproduced (except for 1--)
- hybrid-vector mixing is roughly reproduced
- the constituent picture looks to be a good starting point for detailed modelling; it has the benefit of being a well-constrained, unified description of hadrons and their interactions

# Conclusions

- vector hybrid near 2100 w/ decay constant  $\sim 20$  MeV
- partner states at 2100-2250 and 2220-2350.
- with a  $\pi_1(1600)$  expect partner states at 1750-1780 and  $\sim 1900$  (near the  $\eta_1(1855)$ ).



~thank you~